

Effect of Bypass Fat Supplementation on Post-Partum Reproductive Performance of Crossbred Cows

Tiago Carvalho¹ ●, Jonas Grunewald² ●, Ainhoa Valdivieso³ ●

¹ Scientist, Department of Livestock Production Management, Alentejo Institute of Agrarian Sciences, Beja, Portugal. Email: t.carvalho@aiaes.pt | ORCID: 0000-0001-3394-2764

² Associate Professor, Department of Poultry Science, Allgaeu Institute of Livestock Science, Kempten, Germany. Email: j.grunewald@ails-k.de | ORCID: 0000-0001-0020-1791

³ Assistant Professor, Department of Animal Nutrition, Ebro Valley Institute of Crop Sciences, Tudela, Spain. Email: a.valdivieso@evics.es | ORCID: 0000-0001-3726-1683

ABSTRACT

Rumen-protected, or bypass, fat is widely sold to dairy farmers as a way of helping cows return to cyclicity and conceive sooner after calving, and crossbred cows in warm production systems are often named as likely beneficiaries. This narrative review asks how sound that expectation is. It draws on published work on the chemistry and digestion of protected fats, on energy balance and ovarian activity in early lactation, on the fatty acids proposed to act on follicles, the corpus luteum and the uterus, and on the design and interpretation of supplementation trials, and it reports no new data. Fat supplements can raise the energy density of a ration and may deliver specific fatty acids to reproductive tissues, so a benefit is biologically plausible. Reported fertility responses are nevertheless inconsistent. They seem to depend on the body condition and energy status of the cow, on the fatty acid profile and the timing of supplementation, and on whether intake is protected, and they are measured in trials that are usually small, short and rarely designed around reproductive outcomes. Evidence from crossbred cows specifically is thin, and cost is seldom set against the gain. The review concludes that a modest and conditional benefit is plausible, that its size in crossbred cows is not established, and it sets out what a decisive trial would contain.

Keywords: Bypass fat; Rumen-protected fat; Post-partum anoestrus; Crossbred cows; Energy balance; Fertility; Narrative review

Citation: Carvalho, Grunewald and Valdivieso [2019]. Effect of Bypass Fat Supplementation on Post-Partum Reproductive Performance of Crossbred Cows. DOI: 10.62649/v7.i03.2019.pp29-36

Copyright: © 2019 by the authors. Open access under CC BY 4.0 license.

Article Information: Received: January 12, 2019 Accepted: April 20, 2019 Published: August 20, 2019

Article Class: Review Article

1. Introduction

The weeks after calving decide a good deal of what a dairy cow will earn. Milk production climbs quickly while appetite lags behind, the cow draws on her own body reserves, and in many animals the ovaries stay quiet for longer than the farmer would like [1, 2]. Every extra week before she conceives again lengthens the calving interval and adds to the cost of keeping her. For smallholders

with crossbred cows, who work in warm climates on rations that are often short of energy, the problem is familiar and the margin for error is thin [3, 4].

Bypass fat, also called rumen-protected or rumen-inert fat, is one of the feeding tools offered in response. The idea is straightforward. A fat that survives the rumen adds energy to the ration without upsetting fermentation, and some of the

fatty acids it delivers may act directly on the ovary and uterus [5, 6]. Supplements are marketed on that promise, and advisory leaflets often say that cows given them return to heat sooner and conceive more easily.

What the leaflets rarely say is how uneven the evidence is. Some trials report better fertility after fat supplementation and others report nothing, and a few report something worse [7, 8]. This narrative review asks why, draws on published work only and reports no new data. The aim is a fair account of what is known and what is merely repeated.

2. Literature Review

2.1 What Bypass Fat Is and How It Behaves

Dietary fat can disturb the rumen. Free unsaturated fatty acids coat feed particles and can inhibit the fibre-digesting microbes, so fermentation and intake suffer when too much is fed in an unprotected form [9, 10]. Protected products were designed to avoid this. Calcium salts of fatty acids, prilled and hydrogenated fats and some treated oilseeds stay largely inert in the rumen and release their lipid further down the tract (Table 1). Protection is rarely complete, and the fatty acids that are released in the rumen are subject to biohydrogenation, which turns unsaturated acids into more saturated ones and generates intermediates with effects of their own [11].

Table 1. Common forms of fat offered to dairy cows and how each is handled by the rumen.

Form of fat	How it is protected	Points to note
Calcium salts of fatty acids	Poorly soluble at rumen pH, dissociate lower in the tract [9, 10]	Widely used; dissociation is incomplete under some conditions
Prilled saturated fat	Melting point and saturation limit interaction with microbes [9]	Mostly saturated, so offers little of the unsaturated acids proposed for reproduction
Whole oilseeds	Seed coat slows release; protection is partial [13]	Unsaturated acids partly hydrogenated in the rumen [11]
Free vegetable or fish oil	None	Can disturb fermentation and fibre digestion at higher levels [9, 10]
Chemically treated fat or oilseed	Heat or aldehyde treatment of the carrier	Regulatory and quality questions; check local rules

Note: Qualitative summary of the cited reviews. Products differ between manufacturers, so label claims should be checked against independent data.

These intermediates matter beyond the cow's reproductive tract. Some trans fatty acids formed during incomplete biohydrogenation are known to depress milk fat synthesis [12], which is a reminder that the label on the bag does not describe what reaches the animal. Reviews of the effect of added fat on milk production and composition find that responses depend on the fat source and on the diet it joins [13]. A farmer comparing two products is therefore comparing two different delivery systems, not two names for the same thing. This is one reason the same supplement can look excellent in one report and unremarkable in the next.

2.2 Energy Balance and the Quiet Ovary

The best-supported reason for expecting a benefit from fat is energy. A fresh cow cannot eat enough to match the energy she exports in milk, and the deficit is made up from body fat [2, 14]. The depth and length of that negative energy balance have long been linked to the delay before the first post-partum ovulation, because the pulsatile hormone signals that drive follicle growth are suppressed when the cow is short of energy [1, 15]. Metabolic hormones appear to carry the message from the body to the ovary [16, 17].

Body condition is the visible record of this story. Cows that lose too much condition after calving tend to resume cycles later, and body condition score at calving and at breeding is related to reproductive performance [18, 19]. At the other end, cows that are over-conditioned at drying off often eat poorly after calving and suffer more metabolic trouble [20]. A feed supplement is only one of many overlapping risk factors for late resumption of cycles [21, 22].

If energy is the limiting step, then fat is a sensible tool, since it is the most energy-dense ingredient a farmer can add. The weakness of the argument is that it works only if the cow actually eats the extra energy. High levels of supplemental fat can reduce intake, and a gain in energy density can be cancelled by a fall in the amount eaten [10]. Work with prilled fat in early post-partum dairy cows measured energy balance and follicle development side by side, a design that helps separate an energy effect from a more specific one [23].

2.3 Fatty Acids as Signals

Fatty acids do more than burn. Cholesterol and its precursors supply material for steroid hormone synthesis, and some fatty acids influence the release of prostaglandin from the uterus and the growth of follicles [5, 6]. Work in beef cows found that the lipid composition of the diet changed

luteal activity and follicular growth, and that fats differing in fatty acid composition can act differently even when energy intake is matched [24, 25]. The idea that type matters as much as amount has shaped much of the later research (Table 2).

Table 2. Proposed routes by which dietary fat may influence post-partum reproduction.

Route	Proposed mechanism	Support in cattle
Energy supply	More energy per kilogram of feed may soften the early-lactation energy deficit [1, 15]	Plausible; depends on intake
Steroid precursors	Cholesterol and fatty acids supply material for steroid hormone synthesis [5, 6]	Plausible; indirect
Follicle growth	Fatty acid type may change follicle size and number [24, 25]	Mixed
Uterine signalling	Specific acids may alter prostaglandin release and the uterine environment [5, 34]	Mixed
Oocyte and embryo	Fatty acids in follicular fluid may affect oocyte quality [26, 28]	Early; mostly mechanistic

Note: The last column is the authors' qualitative reading of how well each route is supported in cattle, not a count of studies.

Studies of omega-3 fatty acids add a further layer. Feeding fats of differing n-6 to n-3 ratio changed the fatty acid composition of the ovarian compartments, the follicles and the oocyte in dairy cows [26], and reviews of n-3 acids describe effects on the reproductive and immune systems that are not yet fully worked out [27]. The environment in the follicle seems to link the cow's metabolism with the quality of the oocyte [28], so a feed that alters what circulates may also alter the egg. How far a protected supplement can do this in a cow that is not otherwise in trouble remains open.

2.4 What Beef and Dairy Trials Have Shown

Beef cows supplied many of the earlier trials. Reviews of fat supplementation in beef females and of the nutritional control of beef cow reproduction conclude that responses vary, and that the type of fat and the condition of the cows are part of the explanation [29, 30]. A decade-long review of fat research in beef cattle and sheep likewise treats the fat source and the animal's nutritional state as part of the answer [31]. Calcium-soap trials in post-partum beef cows examined the return to reproductive function [32], and work in *Bos indicus* beef cows tested protected polyunsaturated acids directly [33].

Dairy studies tell a mixed story too. An early review of dietary fat in dairy cattle considered metabolic disorders and reproductive performance together and did not arrive at a simple verdict [8]. Trials that varied the fatty acids fed in the peripartum and breeding periods measured uterine and metabolic responses along with reproduction, which allows the routes to be examined together [34]. Meta-analytic work has since pooled many dairy trials in an effort to explain their differences, and reports that the fertility response depends on the outcome examined, the type of fat and the animals studied [7, 35]. This is a sound reason for caution before promising any single outcome.

2.5 Crossbred Cows and the Environment

Crossbred cows, typically a taurine dairy breed crossed with zebu stock, combine higher milk yield with better tolerance of heat than purebred taurines, but they sit uneasily between two bodies of research. Follicular dynamics in *Bos indicus* cattle differ in several ways from those of *Bos taurus* [3], and most of the fat trials in dairy cattle were done on Holsteins in temperate settings. Whether a response seen in one carries over to the other is a reasonable question and not a settled one.

Heat adds its own difficulty. Hot weather depresses intake, impairs follicle and oocyte function and reduces fertility in dairy cows [4, 36], and lactating cows with high feed intake also clear steroid hormones faster, which lowers their circulating levels [37]. A supplement that raises energy density might help a heat-stressed cow, but the heat that limits her appetite may also limit her response, so a trial that ignores season can mislead in either direction. Small differences of this kind matter little in a single cow but add up across a herd and a year.

3. Materials and Methods

3.1 Scope and Question

The review asks one question in two parts: does supplementing post-partum crossbred cows with bypass fat plausibly improve reproductive performance, and under what conditions would it be expected to? The question was divided into the points a trial would have to settle: what the cows were fed, how the supplement was made and described, how reproductive outcomes were defined, how body condition and season were handled, and how cost was counted. None of these steps is difficult, but each one is easy to skip.

3.2 Literature Assembly

The review draws on peer-reviewed work from four areas: the chemistry and digestion of protected fats; energy balance and ovarian function in early lactation; the action of fatty acids on follicles, the corpus luteum and the uterus; and the design and statistics of supplementation trials. Work on beef cattle and on purebred dairy cows is used for the mechanisms and the pattern of results, not as direct evidence of what happens in crossbred cows.

We did not run a formal systematic search and we do not claim one. This is a narrative synthesis, and the choice of sources reflects the authors' judgement. The review is therefore a better guide to what a good trial should contain than to the size of any particular effect, and it avoids quoting numerical results from individual trials that could not be checked against the primary papers for this article.

3.3 Appraisal Framework

To judge how far a report can be trusted we asked four questions. Were the cows described in terms of body condition, parity and stage? Was the supplement described fully, including the fatty acid profile and the intake achieved? Was the trial large enough for a reproductive outcome, bearing in mind how unreliable small studies are [38, 39]? And was the experimental unit the cow or the group, and did the analysis match [40, 41]? Table 3 lists the sources of variation we looked for.

Table 3. Why supplementation trials disagree and what would help.


Source of variation	Consequence	What to report
Body condition at calving	A thin cow and a fat cow may respond differently [18, 19]	Score at calving and at breeding
Fatty acid profile	Saturated and unsaturated fats act differently [6, 27]	Full profile of the supplement
Intake not recorded	Reduced intake can cancel the energy gain [10]	Dry matter intake by group
Small numbers	Unstable, exaggerated effects [38, 39]	Interval estimates
Pen as unit	Overstated precision [40]	Analyse at the unit fed
Heat and season	Stress can mask a response [4, 36]	Season and housing

Note: Sources of variation are those discussed in the cited literature and are not attributed to any single study.

4. Results

4.1 Where a Protected Fat Acts

A protected fat has several places to succeed or fail, and the literature is easiest to read layer by layer (Figure 1). In the rumen the question is whether protection holds [9, 11]. In the intestine it is whether the cow digests and absorbs what reaches it. In the blood and tissues it is which fatty acids arrive and whether they alter follicle growth, steroid production or uterine function [5, 6]. A trial that reports only conception cannot say which layer failed.



Schematic of the pathway; not data

Figure 1. Where a protected fat acts, layer by layer, and what can interfere (schematic, not data).

Intake sits across all three layers: a high dose that depresses appetite undermines the energy argument entirely [10], and reports that give the amount offered but not the amount eaten leave the reader guessing. Read this way, the absence of an effect in a small trial is not evidence of absence.

4.2 Timing: When in the Post-Partum Period

Timing is the second thing the results turn on. The sequence runs from calving through uterine recovery, the first follicular wave, the first ovulation and the first service to conception (Figure 2). Energy deficit bites hardest early, in the weeks when the first dominant follicle is selected [15, 23], so a supplement started late may arrive after the damage is done. A supplement given before calving, as in studies of the peripartum period, may influence the uterus as well as the ovary [34]. Fat started after that window cannot undo the delay.




Figure 2. Early post-partum timeline and where fat is proposed to act (schematic, not data).

Many trials start supplementation, and report outcomes, at a convenient point. A null result then cannot say whether fat did nothing or was simply offered at the wrong time. Time to first ovulation, first-service conception and calving interval each reflect a different part of the sequence, so one should not stand in for another [21, 22].

4.3 Who Responds: Body Condition and Energy Status

The most consistent theme across the reviews is that response depends on the cow. Thin cows in deep energy deficit have the most to gain from extra energy, while fat cows may respond poorly or eat less of everything [18, 20]. Figure 3 sets out this reasoning as a matrix. It is a statement of hypotheses to be tested, not a summary of results.

	Ration short of energy	Ration adequate in energy
Thin cow, large deficit	Benefit most plausible	Benefit possible (fatty acid type)
Cow in good condition	Benefit possible	Benefit least likely
Over-conditioned cow	Uncertain; intake may fall	Benefit least likely

Hypotheses to be tested; not data

Figure 3. Qualitative matrix of cow state against expected response (hypothesis, not data).

The pattern would explain why pooled results are heterogeneous. If a trial enrolls mostly well-conditioned cows on generous rations, a null result is to be expected. If it enrolls thin cows on poor rations, a benefit may appear, but the result then says little about herds that are better fed [7, 31]. Reports that give only a herd-level mean hide this, and the same supplement can look effective or useless depending on whom it was tested on.

The point is not that a trial is poor, only that its result travels less far than its abstract suggests.

4.4 Cost and Practical Fit

Economic evidence is sparse. Protected fats are generally more expensive per unit of energy than ordinary energy feeds, so a supplement is worth buying only if the fertility gain is large enough, and valuable enough, to pay for it. A gain in fertility that does not cover the price of the supplement is not a gain a farmer would see in the accounts.

A fair cost analysis would record the supplement's price per cow, the length of feeding, the effect on milk and the change in reproductive outcomes at local prices. Trials that report only conception rates cannot answer the farmer's actual question, which is whether the money is better spent here or on better forage, improved shelter in hot weather or closer heat detection [4, 36].

4.5 Quality of the Evidence

The trials share a set of design limitations (Table 3). Conception is a binary outcome that needs large numbers to detect modest differences, yet trials commonly enrol few cows. Small studies produce unstable estimates and tend to overstate effects when they reach significance [38, 39]. Cows housed together share feed and environment, and treating each cow as an independent replicate then inflates the apparent precision [40]. Methods that combine many trials treat the variation between studies as something to be explained, not averaged away [41].

Put plainly, the literature supports a cautious statement. Protected fat can raise the energy density of the ration, and a benefit to post-partum reproduction is biologically plausible, most clearly in thin cows early in lactation. The size of any benefit in crossbred cows is not established by evidence we can cite with confidence, and the best summary is that it is likely to be modest and conditional on body condition, fat type and timing. That is a modest claim, and it is the one the evidence will currently bear.

5. Discussion

5.1 Interpretation and Practical Reading

The main finding is a conditional one. Bypass fat is most likely to help a thin cow, in early lactation, offered a supplement she actually eats, in a season that does not suppress her appetite. It is least likely to help a cow already in good condition on a generous ration. That fits what is known about energy balance and the ovary in general [15, 17], and it implies that the question for a farmer is not

whether bypass fat works but which of his cows are short of what it supplies. Nothing in this reading argues against feeding bypass fat; it argues for measuring what happens when it is fed.

The fatty acid story adds a second conditional. If specific acids act on the follicle or uterus, a mostly saturated prilled fat should not be expected to behave like a supplement rich in the unsaturated acids discussed in the reproductive literature [6, 26]; yet the profile is seldom reported.

Three findings would change this reading. A large, well-designed trial in crossbred cows showing a clear benefit in well-conditioned animals would point to a route other than energy. A trial showing no benefit in thin cows offered a measured and eaten dose would weaken the energy argument. And a full cost analysis showing a net return on typical smallholdings would settle the economic case for those settings.

5.2 Limitations

This is a narrative review and shares the limits of that form. The selection of sources reflects judgement, no formal search or quality scoring was done, and the review quotes no numerical results from individual trials, because these could not be checked against the primary reports for this article. A reader who needs an effect size should consult the original studies, applying the appraisal questions above, or wait for a formal systematic review. The review also leans on beef, purebred dairy and mechanistic work, which are imperfect proxies for crossbred cows, and it uses them to frame hypotheses. Reviews are more useful when they say where the evidence stops than when they smooth the edges over.

5.3 Implications for Trial Design

A trial that could settle the matter would enrol enough cows to detect a modest difference in a binary outcome, balance them for parity, body condition and yield, and stratify by body condition at calving so that responders can be told from non-responders. It would describe the supplement in full, including the fatty acid profile and the protection method, record the dry matter intake of each group, and start feeding at a defined time before or soon after calving. Outcomes would be defined in advance and cover the whole sequence, from first ovulation to calving interval, with cost recorded alongside. Season and housing would be reported, and the group, not the cow, would be the unit of analysis where cows share feed [40, 41].

5.4 Directions for Future Work

Three needs stand out. First, a formal systematic review of the trials in cattle with a transparent search, quality appraisal and, where the data allow, a meta-analysis that treats variation between studies as a result [7, 35]. Second, well-powered trials in crossbred cows stratified by body condition and conducted across seasons. Third, work to establish which fatty acids reach the follicle and uterus when protected supplements are fed to cows kept in warm conditions [4, 28]. These points apply with equal force to any new supplement, not only to fat.

6. Conclusion

Bypass fat is a plausible aid to the post-partum cow, and the case for it rests on sound physiology: it adds energy to a ration at a time of deficit, and some of its fatty acids may act on the ovary and uterus. The evidence, as we can responsibly summarise it, does not yet establish how much it improves reproductive performance in crossbred cows. The reported responses vary with body condition, energy status, fatty acid profile, timing and season, and the trials behind them are usually too small to separate these influences. Claims of large gains deserve caution until they meet the standards described here.

For the practitioner the message is practical rather than discouraging: bypass fat can be tried, preferably in thin early-lactation cows, with the dose eaten recorded and the costs written down. For the researcher, the opportunity is a set of large, well-described and fully costed trials in crossbred cows.

7. Acknowledgements

The authors thank colleagues who commented on earlier drafts of this review.

8. Funding

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

9. Conflict of Interest

The authors declare no conflict of interest.

10. Author Contributions

T.C. conceived the review and wrote the sections on crossbred cow production and trial design. J.G. wrote the sections on the economics of supplementation. A.V. wrote the sections on rumen lipid metabolism and reproductive physiology. All authors revised the manuscript and approved the

final version.

11. Ethics Statement

This is a review of published literature. No animals were used and no ethical approval was required.

12. Data Availability

No new data were generated. All sources are cited in the reference list.

13. References

- [1] Butler, W. R., & Smith, R. D. (1989). Interrelationships between energy balance and postpartum reproductive function in dairy cattle. *Journal of Dairy Science*, 72(3), 767–783. DOI: [10.3168/jds.S0022-0302\(89\)79169-4](https://doi.org/10.3168/jds.S0022-0302(89)79169-4)
- [2] Drackley, J. K. (1999). Biology of dairy cows during the transition period: The final frontier? *Journal of Dairy Science*, 82(11), 2259–2273. DOI: [10.3168/jds.S0022-0302\(99\)75474-3](https://doi.org/10.3168/jds.S0022-0302(99)75474-3)
- [3] Bó, G. A., Baruselli, P. S., & Martinez, M. F. (2003). Pattern and manipulation of follicular development in *Bos indicus* cattle. *Animal Reproduction Science*, 78(3–4), 307–326. DOI: [10.1016/S0378-4320\(03\)00097-6](https://doi.org/10.1016/S0378-4320(03)00097-6)
- [4] Hansen, P. J. (2009). Effects of heat stress on mammalian reproduction. *Philosophical Transactions of the Royal Society B*, 364(1534), 3341–3350. DOI: [10.1098/rstb.2009.0131](https://doi.org/10.1098/rstb.2009.0131)
- [5] Mattos, R., Staples, C. R., & Thatcher, W. W. (2000). Effects of dietary fatty acids on reproduction in ruminants. *Reviews of Reproduction*, 5(1), 38–45. DOI: [10.1530/ror.0.0050038](https://doi.org/10.1530/ror.0.0050038)
- [6] Santos, J. E. P., Bilby, T. R., Thatcher, W. W., Staples, C. R., & Silvestre, F. T. (2008). Long chain fatty acids of diet as factors influencing reproduction in cattle. *Reproduction in Domestic Animals*, 43(Suppl. 2), 23–30. DOI: [10.1111/j.1439-0531.2008.01139.x](https://doi.org/10.1111/j.1439-0531.2008.01139.x)
- [7] Rodney, R. M., Celi, P., Scott, W., Breinhild, K., & Lean, I. J. (2015). Effects of dietary fat on fertility of dairy cattle: A meta-analysis and meta-regression. *Journal of Dairy Science*, 98(8), 5601–5620. DOI: [10.3168/jds.2015-9528](https://doi.org/10.3168/jds.2015-9528)
- [8] Grummer, R. R., & Carroll, D. J. (1991). Effects of dietary fat on metabolic disorders and reproductive performance of dairy cattle. *Journal of Animal Science*, 69(9), 3838–3852. DOI: [10.2527/1991.6993838x](https://doi.org/10.2527/1991.6993838x)
- [9] Jenkins, T. C. (1993). Lipid metabolism in the rumen. *Journal of Dairy Science*, 76(12), 3851–3863. DOI: [10.3168/jds.S0022-0302\(93\)77727-9](https://doi.org/10.3168/jds.S0022-0302(93)77727-9)
- [10] Allen, M. S. (2000). Effects of diet on short-term regulation of feed intake by lactating dairy cattle. *Journal of Dairy Science*, 83(7), 1598–1624. DOI: [10.3168/jds.S0022-0302\(00\)75030-2](https://doi.org/10.3168/jds.S0022-0302(00)75030-2)
- [11] Jenkins, T. C., Wallace, R. J., Moate, P. J., & Mosley, E. E. (2008). Board-invited review: Recent advances in biohydrogenation of unsaturated fatty acids within the rumen microbial ecosystem. *Journal of Animal Science*, 86(2), 397–412. DOI: [10.2527/jas.2007-0588](https://doi.org/10.2527/jas.2007-0588)
- [12] Bauman, D. E., & Griinari, J. M. (2003). Nutritional regulation of milk fat synthesis. *Annual Review of Nutrition*, 23, 203–227. DOI: [10.1146/annurev.nutr.23.011702.073408](https://doi.org/10.1146/annurev.nutr.23.011702.073408)
- [13] Rabiee, A. R., Breinhild, K., Scott, W., Golder, H. M., Block, E., & Lean, I. J. (2012). Effect of fat additions to diets of dairy cattle on milk production and components: A meta-analysis and meta-regression. *Journal of Dairy Science*, 95(6), 3225–3247. DOI: [10.3168/jds.2011-4895](https://doi.org/10.3168/jds.2011-4895)
- [14] Grummer, R. R. (1995). Impact of changes in organic nutrient metabolism on feeding the transition dairy cow. *Journal of Animal Science*, 73(9), 2820–2833. DOI: [10.2527/1995.7392820x](https://doi.org/10.2527/1995.7392820x)
- [15] Butler, W. R. (2003). Energy balance relationships with follicular development, ovulation and fertility in postpartum dairy cows. *Livestock Production Science*, 83(2–3), 211–218. DOI: [10.1016/S0301-6226\(03\)00112-X](https://doi.org/10.1016/S0301-6226(03)00112-X)
- [16] Diskin, M. G., Mackey, D. R., Roche, J. F., & Sreenan, J. M. (2003). Effects of nutrition and metabolic status on circulating hormones and ovarian follicle development in cattle. *Animal Reproduction Science*, 78(3–4), 345–370. DOI: [10.1016/S0378-4320\(03\)00099-X](https://doi.org/10.1016/S0378-4320(03)00099-X)
- [17] Wathes, D. C., Fenwick, M., Cheng, Z., Bourne, N., Llewellyn, S., Morris, D. G., ... Fitzpatrick, R. (2007). Influence of negative energy balance on cyclicity and fertility in the high producing dairy cow. *Theriogenology*, 68(Suppl. 1), S232–S241. DOI: [10.1016/j.theriogenology.2007.04.006](https://doi.org/10.1016/j.theriogenology.2007.04.006)
- [18] Pryce, J. E., Coffey, M. P., & Simm, G. (2001). The relationship between body condition score and reproductive performance. *Journal of Dairy Science*, 84(6), 1508–1515. DOI: [10.3168/jds.S0022-0302\(01\)70184-1](https://doi.org/10.3168/jds.S0022-0302(01)70184-1)
- [19] Roche, J. R., Friggens, N. C., Kay, J. K., Fisher, M. W., Stafford, K. J., & Berry, D. P. (2009). Invited review: Body condition score and its association with dairy cow productivity, health, and welfare. *Journal of Dairy Science*, 92(12), 5769–5801. DOI: [10.3168/jds.2009-2431](https://doi.org/10.3168/jds.2009-2431)
- [20] Rukkwamsuk, T., Kruip, T. A. M., & Wensing, T. (1999). Relationship between overfeeding and overconditioning in the dry period and the problems of high producing dairy cows during the postparturient period. *Veterinary Quarterly*, 21(3), 71–77. DOI: [10.1080/01652176.1999.9694997](https://doi.org/10.1080/01652176.1999.9694997)
- [21] Santos, J. E. P., Rutigliano, H. M., & Sá Filho, M. F. (2009). Risk factors for resumption of postpartum estrous cycles and embryonic survival in lactating dairy cows. *Animal Reproduction Science*, 110(3–4), 207–221. DOI: [10.1016/j.anireprosci.2008.01.014](https://doi.org/10.1016/j.anireprosci.2008.01.014)
- [22] Lucy, M. C. (2001). Reproductive loss in high-producing dairy cattle: Where will it end?

- Journal of Dairy Science*, 84(6), 1277-1293. DOI: [10.3168/jds.S0022-0302\(01\)70158-0](https://doi.org/10.3168/jds.S0022-0302(01)70158-0)
- [23] Beam, S. W., & Butler, W. R. (1997). Energy balance and ovarian follicle development prior to the first ovulation postpartum in dairy cows receiving three levels of dietary fat. *Biology of Reproduction*, 56(1), 133-142. DOI: [10.1095/biolreprod56.1.133](https://doi.org/10.1095/biolreprod56.1.133)
- [24] Williams, G. L. (1989). Modulation of luteal activity in postpartum beef cows through changes in dietary lipid. *Journal of Animal Science*, 67(3), 785-793. DOI: [10.2527/jas1989.673785x](https://doi.org/10.2527/jas1989.673785x)
- [25] Thomas, M. G., Bao, B., & Williams, G. L. (1997). Dietary fats varying in their fatty acid composition differentially influence follicular growth in cows fed isoenergetic diets. *Journal of Animal Science*, 75(9), 2512-2519. DOI: [10.2527/1997.7592512x](https://doi.org/10.2527/1997.7592512x)
- [26] Zachut, M., Dekel, I., Lehrer, H., Arieli, A., Arav, A., Livshitz, L., ... Moallem, U. (2010). Effects of dietary fats differing in n-6:n-3 ratio fed to high-yielding dairy cows on fatty acid composition of ovarian compartments, follicular status, and oocyte quality. *Journal of Dairy Science*, 93(2), 529-545. DOI: [10.3168/jds.2009-2167](https://doi.org/10.3168/jds.2009-2167)
- [27] Moallem, U. (2018). Invited review: Roles of dietary n-3 fatty acids in performance, milk fat composition, and reproductive and immune systems in dairy cattle. *Journal of Dairy Science*, 101(10), 8641-8661. DOI: [10.3168/jds.2018-14772](https://doi.org/10.3168/jds.2018-14772)
- [28] Leroy, J. L. M. R., Rizos, D., Sturmey, R., Bossaert, P., Gutierrez-Adan, A., Van Hoeck, V., ... Bols, P. E. J. (2012). Intrafollicular conditions as a major link between maternal metabolism and oocyte quality: A focus on dairy cow fertility. *Reproduction, Fertility and Development*, 24(1), 1-12. DOI: [10.1071/RD11901](https://doi.org/10.1071/RD11901)
- [29] Funston, R. N. (2004). Fat supplementation and reproduction in beef females. *Journal of Animal Science*, 82(E-Suppl.), E154-E161. DOI: [10.2527/2004.8213_supplE154x](https://doi.org/10.2527/2004.8213_supplE154x)
- [30] Hess, B. W., Lake, S. L., Scholljegerdes, E. J., Weston, T. R., Nayigihugu, V., Molle, J. D. C., & Moss, G. E. (2005). Nutritional controls of beef cow reproduction. *Journal of Animal Science*, 83(13 Suppl.), E90-E106. DOI: [10.2527/2005.8313_supplE90x](https://doi.org/10.2527/2005.8313_supplE90x)
- [31] Hess, B. W., Moss, G. E., & Rule, D. C. (2008). A decade of developments in the area of fat supplementation research with beef cattle and sheep. *Journal of Animal Science*, 86(14 Suppl.), E188-E204. DOI: [10.2527/jas.2007-0546](https://doi.org/10.2527/jas.2007-0546)
- [32] Hightshoe, R. B., Cochran, R. C., Corah, L. R., Kiracofe, G. H., Harmon, D. L., & Perry, R. C. (1991). Effects of calcium soaps of fatty acids on postpartum reproductive function in beef cows. *Journal of Animal Science*, 69(10), 4097-4103. DOI: [10.2527/1991.69104097x](https://doi.org/10.2527/1991.69104097x)
- [33] Lopes, C. N., Scarpa, A. B., Cappellozza, B. I., Cooke, R. F., & Vasconcelos, J. L. M. (2009). Effects of rumen-protected polyunsaturated fatty acid supplementation on reproductive performance of *Bos indicus* beef cows. *Journal of Animal Science*, 87(12), 3935-3943. DOI: [10.2527/jas.2009-2201](https://doi.org/10.2527/jas.2009-2201)
- [34] Silvestre, F. T., Carvalho, T. S. M., Francisco, N., Santos, J. E. P., Staples, C. R., Jenkins, T. C., & Thatcher, W. W. (2011). Effects of differential supplementation of fatty acids during the peripartum and breeding periods of Holstein cows: I. Uterine and metabolic responses, reproduction, and lactation. *Journal of Dairy Science*, 94(1), 189-204. DOI: [10.3168/jds.2010-3370](https://doi.org/10.3168/jds.2010-3370)
- [35] Rodney, R. M., Celi, P., Scott, W., Breinhild, K., Santos, J. E. P., & Lean, I. J. (2018). Effects of nutrition on the fertility of lactating dairy cattle. *Journal of Dairy Science*, 101(6), 5115-5133. DOI: [10.3168/jds.2017-14064](https://doi.org/10.3168/jds.2017-14064)
- [36] De Rensis, F., & Scaramuzzi, R. J. (2003). Heat stress and seasonal effects on reproduction in the dairy cow: A review. *Theriogenology*, 60(6), 1139-1151. DOI: [10.1016/S0093-691X\(03\)00126-2](https://doi.org/10.1016/S0093-691X(03)00126-2)
- [37] Wiltbank, M., Lopez, H., Sartori, R., Sangsritavong, S., & Gümen, A. (2006). Changes in reproductive physiology of lactating dairy cows due to elevated steroid metabolism. *Theriogenology*, 65(1), 17-29. DOI: [10.1016/j.theriogenology.2005.10.003](https://doi.org/10.1016/j.theriogenology.2005.10.003)
- [38] Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability of neuroscience. *Nature Reviews Neuroscience*, 14(5), 365-376. DOI: [10.1038/nrn3475](https://doi.org/10.1038/nrn3475)
- [39] Ioannidis, J. P. A. (2005). Why most published research findings are false. *PLoS Medicine*, 2(8), e124. DOI: [10.1371/journal.pmed.0020124](https://doi.org/10.1371/journal.pmed.0020124)
- [40] Hurlbert, S. H. (1984). Pseudoreplication and the design of ecological field experiments. *Ecological Monographs*, 54(2), 187-211. DOI: [10.2307/1942661](https://doi.org/10.2307/1942661)
- [41] St-Pierre, N. R. (2001). Invited review: Integrating quantitative findings from multiple studies using mixed model methodology. *Journal of Dairy Science*, 84(4), 741-755. DOI: [10.3168/jds.S0022-0302\(01\)74530-4](https://doi.org/10.3168/jds.S0022-0302(01)74530-4)